

Uncertainty

CHAPTER 14

Outline

- ◇ Uncertainty
- ◇ Probability
- ◇ Syntax
- ◇ Semantics
- ◇ Inference rules

Uncertainty

Let action $A_t =$ leave for airport t minutes before flight
Will A_t get me there on time?

Problems:

- 1) partial observability (road state, other drivers' plans, etc.)
- 2) noisy sensors (KCBS traffic reports)
- 3) uncertainty in action outcomes (flat tire, etc.)
- 4) immense complexity of modelling and predicting traffic

Hence a purely logical approach either

- 1) risks falsehood: " A_{25} will get me there on time"
- or 2) leads to conclusions that are too weak for decision making:
" A_{25} will get me there on time if there's no accident on the bridge
and it doesn't rain and my tires remain intact etc etc."

(A_{1440} might reasonably be said to get me there on time
but I'd have to stay overnight in the airport ...)

Methods for handling uncertainty

Default or nonmonotonic logic:

Assume my car does not have a flat tire

Assume A_{25} works unless contradicted by evidence

Issues: What assumptions are reasonable? How to handle contradiction?

Rules with fudge factors:

$A_{25} \mapsto_{0.3}$ get there on time

$Sprinkler \mapsto_{0.99} WetGrass$

$WetGrass \mapsto_{0.7} Rain$

Issues: Problems with combination, e.g., *Sprinkler* causes *Rain*??

Probability

Given the available evidence,

A_{25} will get me there on time with probability 0.04

Mahaviracarya (9th C.), Cardano (1565) theory of gambling

(Fuzzy logic handles *degree of truth* NOT uncertainty e.g.,
 $WetGrass$ is true to degree 0.2)

Probability

Probabilistic assertions *summarize* effects of
laziness: failure to enumerate exceptions, qualifications, etc.
ignorance: lack of relevant facts, initial conditions, etc.

Subjective or Bayesian probability:

Probabilities relate propositions to one's own state of knowledge
e.g., $P(A_{25} | \text{no reported accidents}) = 0.06$

These are not assertions about the world

Probabilities of propositions change with new evidence:

e.g., $P(A_{25} | \text{no reported accidents, 5 a.m.}) = 0.15$

(Analogous to logical entailment status $KB \models \alpha$, not truth.)

Making decisions under uncertainty

Suppose I believe the following:

$$P(A_{25} \text{ gets me there on time} | \dots) = 0.04$$

$$P(A_{90} \text{ gets me there on time} | \dots) = 0.70$$

$$P(A_{120} \text{ gets me there on time} | \dots) = 0.95$$

$$P(A_{1440} \text{ gets me there on time} | \dots) = 0.9999$$

Which action to choose?

Depends on my preferences for missing flight vs. airport cuisine, etc.

Utility theory is used to represent and infer preferences


Decision theory = utility theory + probability theory

Axioms of probability

For any propositions A, B

1. $0 \leq P(A) \leq 1$
2. $P(\text{True}) = 1$ and $P(\text{False}) = 0$
3. $P(A \vee B) = P(A) + P(B) - P(A \wedge B)$

True



de Finetti (1931): an agent who bets according to probabilities that violate these axioms can be forced to bet so as to lose money regardless of outcome.

Syntax

Similar to propositional logic: possible worlds defined by assignment of values to random variables.

Propositional or Boolean random variables

e.g., *Cavity* (do I have a cavity?)

Include propositional logic expressions

e.g., $\neg \textit{Burglary} \vee \textit{Earthquake}$

Multivalued random variables

e.g., *Weather* is one of $\langle \textit{sunny}, \textit{rain}, \textit{cloudy}, \textit{snow} \rangle$

Values must be exhaustive and mutually exclusive

Proposition constructed by assignment of a value:

e.g., $\textit{Weather} = \textit{sunny}$; also $\textit{Cavity} = \textit{true}$ for clarity

Syntax contd.

Prior or unconditional probabilities of propositions

e.g., $P(Cavity) = 0.1$ and $P(Weather = sunny) = 0.72$
correspond to belief prior to arrival of any (new) evidence

Probability distribution gives values for all possible assignments:

$\mathbf{P}(Weather) = \langle 0.72, 0.1, 0.08, 0.1 \rangle$ (normalized, i.e., sums to 1)

Joint probability distribution for a set of variables gives values for each possible assignment to all the variables

$\mathbf{P}(Weather, Cavity) =$ a 4×2 matrix of values:

<i>Weather =</i>	<i>sunny rain cloudy snow</i>
<i>Cavity = true</i>	
<i>Cavity = false</i>	

Syntax contd.

Conditional or posterior probabilities

e.g., $P(\textit{Cavity}|\textit{Toothache}) = 0.8$

i.e., given that *Toothache* is all I know

Notation for conditional distributions:

$\mathbf{P}(\textit{Weather}|\textit{Earthquake}) = 2\text{-element vector of 4-element vectors}$

If we know more, e.g., *Cavity* is also given, then we have

$P(\textit{Cavity}|\textit{Toothache}, \textit{Cavity}) = 1$

Note: the less specific belief *remains valid* after more evidence arrives, but is not always *useful*

New evidence may be irrelevant, allowing simplification, e.g.,

$P(\textit{Cavity}|\textit{Toothache}, \textit{49ersWin}) = P(\textit{Cavity}|\textit{Toothache}) = 0.8$

This kind of inference, sanctioned by domain knowledge, is crucial

Conditional probability

Definition of conditional probability:

$$P(A|B) = \frac{P(A \wedge B)}{P(B)} \text{ if } P(B) \neq 0$$

Product rule gives an alternative formulation:

$$P(A \wedge B) = P(A|B)P(B) = P(B|A)P(A)$$

A general version holds for whole distributions, e.g.,

$$\mathbf{P}(Weather, Cavity) = \mathbf{P}(Weather|Cavity)\mathbf{P}(Cavity)$$

(View as a 4×2 set of equations, *not* matrix mult.)

Chain rule is derived by successive application of product rule:

$$\begin{aligned} \mathbf{P}(X_1, \dots, X_n) &= \mathbf{P}(X_1, \dots, X_{n-1}) \mathbf{P}(X_n|X_1, \dots, X_{n-1}) \\ &= \mathbf{P}(X_1, \dots, X_{n-2}) \mathbf{P}(X_{n-1}|X_1, \dots, X_{n-2}) \mathbf{P}(X_n|X_1, \dots, X_{n-1}) \\ &= \dots \\ &= \prod_{i=1}^n \mathbf{P}(X_i|X_1, \dots, X_{i-1}) \end{aligned}$$

Bayes' Rule

Product rule $P(A \wedge B) = P(A|B)P(B) = P(B|A)P(A)$

$$\Rightarrow \text{Bayes' rule } P(A|B) = \frac{P(B|A)P(A)}{P(B)}$$

Why is this useful???

For assessing diagnostic probability from causal probability:

$$P(Cause|Effect) = \frac{P(Effect|Cause)P(Cause)}{P(Effect)}$$

E.g., let M be meningitis, S be stiff neck:

$$P(M|S) = \frac{P(S|M)P(M)}{P(S)} = \frac{0.8 \times 0.0001}{0.1} = 0.0008$$

Note: posterior probability of meningitis still very small!

Normalization

Suppose we wish to compute a posterior distribution over A given $B = b$, and suppose A has possible values $a_1 \dots a_m$

We can apply Bayes' rule for each value of A :

$$P(A = a_1 | B = b) = P(B = b | A = a_1)P(A = a_1) / P(B = b)$$

...

$$P(A = a_m | B = b) = P(B = b | A = a_m)P(A = a_m) / P(B = b)$$

Adding these up, and noting that $\sum_i P(A = a_i | B = b) = 1$:

$$1 / P(B = b) = 1 / \sum_i P(B = b | A = a_i)P(A = a_i)$$

This is the normalization factor, constant w.r.t. i , denoted α :

$$\mathbf{P}(A | B = b) = \alpha \mathbf{P}(B = b | A) \mathbf{P}(A)$$

Typically compute an unnormalized distribution, normalize at end

e.g., suppose $\mathbf{P}(B = b | A) \mathbf{P}(A) = \langle 0.4, 0.2, 0.2 \rangle$

then $\mathbf{P}(A | B = b) = \alpha \langle 0.4, 0.2, 0.2 \rangle = \frac{\langle 0.4, 0.2, 0.2 \rangle}{0.4 + 0.2 + 0.2} = \langle 0.5, 0.25, 0.25 \rangle$

Conditioning

Introducing a variable as an extra condition:

$$P(X|Y) = \sum_z P(X|Y, Z = z)P(Z = z|Y)$$

Intuition: often easier to assess each specific circumstance, e.g.,

$$\begin{aligned} P(\text{RunOver}|\text{Cross}) \\ &= P(\text{RunOver}|\text{Cross}, \text{Light} = \text{green})P(\text{Light} = \text{green}|\text{Cross}) \\ &+ P(\text{RunOver}|\text{Cross}, \text{Light} = \text{yellow})P(\text{Light} = \text{yellow}|\text{Cross}) \\ &+ P(\text{RunOver}|\text{Cross}, \text{Light} = \text{red})P(\text{Light} = \text{red}|\text{Cross}) \end{aligned}$$

When Y is absent, we have summing out or marginalization:

$$P(X) = \sum_z P(X|Z = z)P(Z = z) = \sum_z P(X, Z = z)$$

In general, given a joint distribution over a set of variables, the distribution over any subset (called a marginal distribution for historical reasons) can be calculated by summing out the other variables.

Full joint distributions

A complete probability model specifies every entry in the joint distribution for all the variables $\mathbf{X} = X_1, \dots, X_n$

I.e., a probability for each possible world $X_1 = x_1, \dots, X_n = x_n$

(Cf. complete theories in logic.)

E.g., suppose *Toothache* and *Cavity* are the random variables:

	<i>Toothache = true</i>	<i>Toothache = false</i>
<i>Cavity = true</i>	0.04	0.06
<i>Cavity = false</i>	0.01	0.89

Possible worlds are mutually exclusive $\Rightarrow P(w_1 \wedge w_2) = 0$

Possible worlds are exhaustive $\Rightarrow w_1 \vee \dots \vee w_n$ is *True*

hence $\sum_i P(w_i) = 1$

Full joint distributions contd.

- 1) For any proposition ϕ defined on the random variables $\phi(w_i)$ is true or false
- 2) ϕ is equivalent to the disjunction of w_i s where $\phi(w_i)$ is true

$$\text{Hence } P(\phi) = \sum_{\{w_i: \phi(w_i)\}} P(w_i)$$

I.e., the unconditional probability of any proposition is computable as the sum of entries from the full joint distribution

Conditional probabilities can be computed in the same way as a ratio:

$$P(\phi|\xi) = \frac{P(\phi \wedge \xi)}{P(\xi)}$$

E.g.,

$$P(\text{Cavity}|\text{Toothache}) = \frac{P(\text{Cavity} \wedge \text{Toothache})}{P(\text{Toothache})} = \frac{0.04}{0.04 + 0.01} = 0.8$$

Inference from joint distributions

Typically, we are interested in

the posterior joint distribution of the query variables \mathbf{Y}
given specific values \mathbf{e} for the evidence variables \mathbf{E}

Let the hidden variables be $\mathbf{H} = \mathbf{X} - \mathbf{Y} - \mathbf{E}$

Then the required summation of joint entries is done by summing out the hidden variables:

$$\mathbf{P}(\mathbf{Y}|\mathbf{E} = \mathbf{e}) = \alpha \mathbf{P}(\mathbf{Y}, \mathbf{E} = \mathbf{e}) = \alpha \sum_{\mathbf{h}} \mathbf{P}(\mathbf{Y}, \mathbf{E} = \mathbf{e}, \mathbf{H} = \mathbf{h})$$

The terms in the summation are joint entries because \mathbf{Y} , \mathbf{E} , and \mathbf{H} together exhaust the set of random variables

Obvious problems:

- 1) Worst-case time complexity $O(d^n)$ where d is the largest arity
- 2) Space complexity $O(d^n)$ to store the joint distribution
- 3) How to find the numbers for $O(d^n)$ entries???